



DBT Treatment Research

December 1, 2016-January 26, 2017

1. **Title:** [The effect of therapist use of validation strategies on change in client emotion in individual dbt treatment sessions.](#)

Authors: A Carson-Wong, CD Hughes, SL Rizvi

Journal & Date: Personality Disorders (2016)

Dialectical behavior therapy (DBT) is a treatment for borderline personality disorder, a disorder for which emotion dysregulation is central. Within DBT, there are 6 explicitly defined validation strategies that range hierarchically from validation level (VL) 1 to VL 6. To date, there have been no studies on the frequency of use of VLs in actual DBT sessions. The aim of the current study was to explore DBT therapists' use of VLs and examine the relationship between VLs and change in a client emotion during therapy sessions. DBT treatment sessions (n = 121) across 35 participants in a DBT training clinic were coded for therapist use of VLs. A repeated-measures analysis of variance (ANOVA) was used to assess for change in therapist use of VLs over time and hierarchical linear modeling was used to correlate therapist use of these strategies with change in client emotion. Results indicated no significant relationship between overall frequency of VLs and change in client emotion. However, an increase in frequency of high VLs was associated with an increase in positive affect (PA) and a decrease in negative affect (NA) while an increase in frequency of low VLs was associated with a decrease in PA and no change in NA. An increase in frequency of VL 4 was associated with an increase in NA. VL 6 was associated with an increase in PA and a decrease in NA. Findings suggest that specific validation strategies may be related to session changes in affect and have implications for identifying potential mechanisms of change.

Dialectical behavior therapy (DBT) is a treatment for borderline personality disorder, a disorder for which emotion dysregulation is central. Within DBT, there are 6 explicitly defined validation strategies that range hierarchically from validation level (VL) 1 to VL 6. To date, there have been no studies on the frequency of use of VLs in actual DBT sessions. The aim of the current study was to explore DBT therapists' use of VLs and examine the relationship between VLs and change in a client emotion during therapy sessions. DBT treatment sessions (n = 121) across 35 participants in a DBT training clinic were coded for therapist use of VLs. A repeated-measures analysis of variance (ANOVA) was used to assess for change in therapist use of VLs over time and hierarchical linear modeling was used to correlate therapist use of these strategies with change in client emotion. Results indicated no significant relationship between overall frequency of VLs and change in client emotion. However, an increase in frequency of high VLs was associated with an increase in positive affect (PA) and a decrease in negative affect (NA) while an increase in frequency of low VLs was associated with a decrease in PA and no change in NA. An increase in frequency of VL 4 was associated with an increase in NA. VL 6 was associated with an increase



in PA and a decrease in NA. Findings suggest that specific validation strategies may be related to session changes in affect and have implications for identifying potential mechanisms of change.

2.

Title: [Dialectical Behavior Therapy Group Skills Training for Bipolar Disorder](#)

Authors: L Eisner, D Eddie, R Harley, M Jacobo, AA Nierenberg, T Deckersbach

Journal & Date: Behavior Therapy (2017)

There is growing evidence that the capacity for emotion regulation is compromised in individuals with bipolar disorder. Dialectical behavior therapy (DBT), an empirically supported treatment that specifically targets emotion dysregulation, may be an effective adjunct treatment for improving emotion regulation and residual mood symptoms in patients with bipolar disorder. In this open, proof of concept pilot study, 37 participants engaged in a 12-week DBT group skills training program, learning mindfulness, emotion regulation, and distress tolerance skills. Repeated measures mixed models revealed skill acquisition in the areas of mindfulness, emotion regulation and distress tolerance, as well as improved psychological well-being and decreased emotion reactivity. The results of this study support a burgeoning literature that DBT is a feasible adjunct intervention for patients with bipolar disorder.

3.

Title: [Treating Depression Among Adolescent Perinatal Women With a Dialectical Behavior Therapy-Informed Skills Group](#)

Authors: BV Kleiber, JN Felder, B Ashby, S Scott, J Dean, S Dimidjian

Journal & Date: Cognitive and Behavioral Practice (2017)

Depression is a prevalent and impairing problem affecting both women and offspring during the perinatal (pregnancy and the postpartum) period. Despite this, few studies have examined treatments for perinatal adolescents with depressive symptoms. The present study examined the feasibility and preliminary outcomes of a Dialectical Behavioral Therapy (DBT) informed skills group among depressed adolescent perinatal women (N = 25) recruited from both a public health parent education program and an adolescent obstetric clinic. A brief composite case example is included to illustrate how DBT skills were taught, practiced, and applied. Findings suggest the intervention was credible, acceptable, and associated with improvement in depression. Challenges with feasibility of enrolling and retaining adolescent perinatal women were evident, as approximately half of the enrolled participants did not complete the study. This study provides preliminary evidence that a DBT-informed skills group may be a promising intervention for depressed adolescent perinatal women and points to important directions for clinical practice and research, including treatment engagement and retention.

4. **Title:** [Experiences of women with learning disabilities undergoing dialectical behaviour therapy in a secure service](#)

Authors: M Thomson, P Johnson

Journal & Date: British Journal of Learning Disabilities (2017)



Background: This research aimed to capture the experiences of women with learning disabilities living in secure services who undertook dialectical behaviour therapy (DBT); they were the first people to do this in the trust. It is hoped their experiences may guide and inform other services undertaking the same process.

Materials and Methods: Interpretative phenomenological analysis (IPA) was used to provide a flexible framework to make sense of the way the women interpreted their experiences.

Results and Discussion: The analysis identified three main themes: How you do DBT, What we think about DBT, Using DBT. The qualitative approach adds a valuable contribution to the wider literature, highlighting the importance of capturing the women's voices.

Conclusion: As most published literature regarding the effectiveness of DBT is not written from first hand service user experiences, it is hoped this research offers a unique insight into these women's experience.

5. **Title:** [Exploring the effectiveness of combined mentalization-based group therapy and dialectical behaviour therapy for inpatients with borderline personality disorder – A pilot study](#)

Authors: MA Edel, V Raaff, G Dimaggio, A Buchheim, M Brune

Journal & Date: British Journal of Clinical Psychology (2016)

Objectives: Borderline personality disorder (BPD) is characterized by emotional instability, interpersonal dysfunction, and other features that typically develop before a background of insecure attachment and traumatic experiences. Dialectical behaviour therapy (DBT) has proven highly effective in reducing self-harm and improving emotion regulation, whereby problems concerning social cognition, which are also characteristic of BPD, may need additional approaches such as mentalization-based treatment (MBT).

Methods: Here, we examined, in a pilot study, the effectiveness of MBT given adjunct to DBT, compared to DBT alone, in an inpatient sample with BPD, whereby mentalization was measured using a novel cartoon-based task.

Results: Both treatments were highly effective in reducing symptom severity. The combination of DBT and MBT was superior in reducing fearful attachment and in improving affective mentalizing.

Conclusions: Mentalization-based treatment in combination with DBT may improve certain aspects of social cognitive skills and attachment security, as compared to DBT alone, although the exact mechanisms that led to these changes need to be studied further.